



# DCM LEAFGEL® NK

## Foliar fertiliser NK 8-4 gel formulation with plant amino acids

### Composition

Fertiliser – Suspension of organo-mineral compound fertiliser NK with plant amino acids

- 8% TOTAL NITROGEN (N) of which:
  - 8% organically-bound nitrogen (from plant amino acids)
- 4% POTASSIUM OXIDE (K<sub>2</sub>O) soluble in water
- 40% TOTAL AMINO ACIDS

Product authorised for use in organic agriculture in accordance with Regulation (EU) 2018/848 and its amendments.



### Characteristics

- foliar fertiliser with a high percentage of amino acids, which are 100% of plant origin
- contains no ingredients of animal origin
- the amino acids have been specifically selected because they are easily uptakeable by the leaves, which ensures a rapid effect on growth and colour
- in addition to amino acids, the products contains other components that facilitate uptake into the leaves
- prevents and remedies deficiency symptoms
- almost odourless, light brown gel
- **with biostimulant effect:** the amino acids ensure a strong crop that is more resistant to abiotic stress factors
- can be used both continuously and periodically for adjustments during crop growth
- the uptake is independent of the soil conditions (pH, soil temperature)
- the strictly controlled production process ensures a high purity and permanently stable composition

### Form

**gel formulation** (suspension):

- highly concentrated product
- easy dosing
- highly water-soluble
- remains stable (no precipitation)
- can be applied in combination with other products

density: 1400 g/L

### Packaging

5 kg container (4 containers per carton)





# DCM LEAFGEL® NK

**Foliar fertiliser NK 8-4 gel formulation  
with amino acids of plant origin**

## Instructions for use

Foliar fertilization: ..... 2 – 3 gram/L solution  
At least 2 applications, with an interval of 10 days

The exact dose depends on the needs of the crop, the time of application (growth stage), the nutrient reserve in the soil/substrate and the intensity of irrigation.

Apply in the active growing season (spring - summer), as soon as the plants are well in leaf. Apply preferably in the late afternoon or early evening, out of the sun, on slightly moist plants and before the opening of the flower buds.

Shake the container well before use. Make the correct dilution with water and fertiliser and spray the solution evenly over the leaves until it drips down from the plants (drip-off).

Contact your dealer for additional product compatibility information.

## Storage

Close packaging securely after use.

Store at a temperature between 5 and 25 °C.

Keep out of reach of children and pets.

DCM products meet the nutritional values indicated on their packaging and/or the technical data sheets, and are fully traceable. Product advice is intended to be used for informative purposes only and does not imply any commitment or agreement. The instructions for use are based on many years of practical experience and research. Each plant and each cropping system has its own nutritional requirements. The time of application, nutrient reserves in the soil/growing medium and statutory fertilization standards are also important in determining fertiliser application rates. It is advisable (good practice) to always test new product applications on a small scale first. Fertilisers may cause EC to increase and can affect the pH level. All of these factors are essential to consider when using one fertiliser product or when combining different fertilisers. Growing media to which fertiliser products have been added should be used as soon as possible after delivery. DCM accepts no liability for consequential damage caused by the use of its products.

TECHNICAL DATA SHEET FOR EXPORT – EXEN–KVE-240829