



DCM OLEGA® FER

3 % N + 3 % Fe + seaweed extract

Composition

Solution of fertilizers with iron and seaweed extract
for foliar fertilization

- 3 % TOTAL NITROGEN (N) of which
3 % ureic nitrogen
- 3 % IRON (Fe) soluble in water, of which
2 % iron (Fe) complexed by citrate
1 % iron (Fe) DTPA chelated
- 3 % SEAWEED EXTRACT



Characteristics

- **foliar fertilizer** with plant own iron (Fe) complex and seaweed extract
 - faster and better iron release when sprayed onto the leaves
 - better uptake through the leaves due to the small molecule
 - added urea for an improved uptake of iron (synergy)
 - quicker translocation in the xylem and incorporation in the plant
- faster and more intense greening response
- prevents and remedies iron deficiency
(symptoms: chlorosis of the youngest leaves, growth inhibition, potential leaf fall)
- lower risk of leaf scorching
- quicker uptake than soil fertilization and irrespective of soil conditions (pH, soil temperature)
- easily applicable

Form

suspension – liquid

Packaging

drums of 15 L (16,5 kg)
drums of 5 L (5,5 kg) – 4 drums/package

Storage

Store in a frost-free place.





DCM OLEGA[®] FER

3 % N + 3 % Fe + seaweed extract

Instructions for use

- To be used in the active growing season (spring – summer), as soon as the plants are well in leaf.
- Apply preferably in the late afternoon or early evening, on slightly moist plants.
- Do not apply in the blazing sun and at temperatures above 25°C.
- Do not apply on opened flowers.
- Shake the bottle well before use.
- A foliar fertiliser is to be used supplementary to soil fertilization, not as a substitute.
- Preferably use preventively in order to avoid growth inhibition due to iron deficiency.
- Effect and quickness of response are dependent on the plant, the variety, the growth stage and the climatic circumstances: therefore always use the lowest dosage during the first application.
- In combination with a wetting agent, do not exceed the concentration of 10 ml per litre of water.

Dosage

The exact rate depends on the soil conditions, the nutrient reserves in the soil and the needs of the crop. Ask for our specialized advice.

FRUIT TREES (f.i. PEARS), PERENNIAL, WOODY SHRUBS AND TREES (before opening of the buds)

- preventive 1,5 - 7 L/ha
- curative 4 - 7 L/ha

always dilute with minimum 400 litres of water per hectare

according to necessity, minimum 2, preferably 3 applications with an interval of 10 – 14 days

ANNUAL PLANTS AND FLOWERS (before opening of the buds), VEGETABLES AND HERBS

- preventive 1 - 4 L/ha
- curative 2 - 4 L/ha
- crops under glass max. 2,4 L/ha

always dilute with minimum 400 litres of water per hectare

according to necessity, minimum 2, preferably 3 applications with an interval of 10 – 14 days

LAWNS AND SPORTS FIELDS

- standard dosage 7 - 10 L/ha
always dilute with minimum 400 litres of water per hectare
according to necessity, minimum 3, preferably 5 applications with an interval of 10 – 14 days
- fast greening response 40 - 60 L/ha
dilute with a minimum of 1000 litres of water per hectare

APPLICATIONS IN THE LAYING-OUT OF GARDENS AND ON SMALL SURFACES

- perennial, woody shrubs and trees (before opening of the buds) 8 - 15 ml/L water
- annual plants and flowers (before opening of the buds) 4 - 6 ml/L water
make the correct dilution with water and spray evenly over the leaves until it drips down from the plants

DCM products meet the nutritional values indicated on their packaging and/or the technical data sheets, and are fully traceable. Product advice is intended to be used for informative purposes only and does not imply any commitment or agreement. The instructions for use are based on many years of practical experience and research. Each plant and each cropping system has its own nutritional requirements. The time of application, nutrient reserves in the soil/growing medium and statutory fertilization standards are also important in determining fertiliser application rates. It is advisable (good practice) to always test new product applications on a small scale first. Fertilisers may cause EC to increase and can affect the pH level. All of these factors are essential to consider when using one fertiliser product or when combining different fertilisers. Growing media to which fertiliser products have been added should be used as soon as possible after delivery. DCM accepts no liability for consequential damage caused by the use of its products.

TECHNICAL DATA SHEET FOR EXPORT – EXEN – DBOE - 220331